Author Index

Alexander, R., see Forster, P. G. 384
Ansari, G. A. S., see Bhat, H. K. 369
Armstrong, D. W., see Seeman, J. I. 169
Ayyangar, N. R.
——, Tambe, A. S. and Biswas, S. S.
Evaluation of 4-(2′-phenylisopropyl)phenol esters as stationary phases for gas–liquid chromatography 33
Beenackers, A. A. C. M., see Marsman, J. H. 111
Bhat, H. K.
—— and Ansari, G. A. S.
Improved separation of lipid esters by thin-layer chromatography 369
Bingham, B., see Newland, P. 406
Biswas, S. S., see Ayyangar, N. R. 33
Blessington, B.
—— and Crabb, N.
Proposed primary reference methods for the determination of some commercially important chiral aryloxypropionate herbicides in both free acid and ester forms 349
Bogart, G. R.
——, Leyden, D. E., Wade, T. M., Schafer, W. and Carr, P. W.
Spectroscopic investigation of the hydrolysis of γ-glycidoxypropylsilane bound to silica surfaces 209
Braumann, T., see Techel, D. 427
Brown, J. M., see Whitley, R. D. 263
Brückner, C., see Schneider, G. 459
Cadelis, F., see Kirchberger, J. 289
Carr, P. W., see Bogart, G. R. 209
Casoli, A.
——, Mangia, A., Predieri, G. and Sappa, E.
High-performance liquid chromatography of triangular ruthenium clusters containing bidentate group-15 donor ligands 443
Chiba, M., see Vukmanic, D. 189
Collins, D. A.
Modification to a thermal energy analyzer with associated electronic filtering for improved gas chromatographic analysis of explosives traces 379
Cox, G. B.
—— and Snyder, L. R.
Preparative high-performance liquid chromatography under isocratic conditions. III. The consequences of two adjacent bands having unequal column capacities 95
——, see Snyder, L. R. 63, 85
Crabb, N., see Blessington, B. 349
Davies, D. S., see Williams, J. A. 454
Davies, P. A.
Determination of diffusion coefficients of proteins in beaded agarose by gel filtration 221
Demarco, A. C., see White, R. L. 437
Dent, M. R., see Price, G. J. 1
Dolan, J. W., see Snyder, L. R. 63
Dratz, E. A., see Miercke, L. J. W. 331
Euerby, M. R.
——, Partridge, L. Z. and Gibbons, W. A.
Study of the chromatographic behaviour and resolution of α-amino acid enantiomers by high-performance liquid chromatography utilizing pre-column derivatization with *o*-phthaldialdehyde and new chiral thiols 239
Ford, J. C.
—— and Smith, J. A.
Synthetic peptide purification by application of linear solvent strength gradient theory 131
Forster, P. G.
——, Alexander, R. and Kagi, R. I.
Identification and analysis of tetramethylnaphthalenes in petroleum 384
Frankena, H., see Schotman, P. 301
Fu, R., see Jin, Y. 394
Gibbons, W. A., see Euerby, M. R. 239
Gierak, A.
—— and Leboda, R.
Preparation of carboneous adsorbents by catalytic decomposition of hydrocarbons for chromatographic application 197
Gispen, W. H., see Schotman, P. 301
Goddijn, O. J. M., see Pennings, E. J. M. 311
Gong, J., see Jin, Y. 394
Guan, Z., see Jin, Y. 394
Haasnoot, W.
——, Stouten, P. and Venema, D. P.
High-performance liquid chromatography determination of the proteolysis in Gouda cheese 319
Hancock, F. E., see Smith, J. R. L. 341
Hashimoto, T., see Kato, Y. 401
Herraiz, M., see Reglero, G. 43
Herraiz, T., see Reglero, G. 43
Hoge, J. H. C., see Pennings, E. J. M. 311
Huang, S.-D., see Wang, W.-K. 121
Hudgens, E. E., see Smith, D. F. 431
Jin, Y.
——, Fu, R., Guan, Z., Gong, J. and Li, B.
Low-temperature side-chain liquid crystalline polysiloxanes used as stationary phases for capillary gas chromatography 394
Kagi, R. I., see Forster, P. G. 384

Kakihana, H.
—— and Oi, T.
Theory of isotope separation by displacement chromatography 179
Kamounah, F. S., see Salman, S. R. 390
Karajgikar, N. P., see Whitley, R. D. 263
Kato, Y.
——, Kitamura, T., Nakatani, S. and Hashimoto, T.
High-performance hydrophobic interaction chromatography of proteins on a pellicular support based on hydrophilic resin 401
Kemmelmeier, C., see Paterson, R. R. M. 153
Kim, L.-S.
——, Tian, Z.-Q., Li, X.-N., Zhang, Y. and Wang, J.
Centrifugal counter-current partition chromatography with helical coil rotor. Simplified counter-current chromatography with a rotating face-seal 359
Kirchberger, J.
——, Cadelis, F., Kopperschläger, G. and Vijayalakshmi, M. A.
Interaction of lactate dehydrogenase with structurally related triazine dyes using affinity partitioning and affinity chromatography 289
Kitamura, T., see Kato, Y. 401
Kleindienst, T. E., see Smith, D. F. 431
Kopperschläger, G., see Kirchberger, J. 289
Kramell, R., see Schneider, G. 459
Landrum, D. C.
—— and Mawhinney, T. P.
Gas–liquid chromatography–mass spectrometry of mono- and dithiols as their *tert.*-butyldimethylsilyl derivatives 21
Leboda, R., see Gierak, A. 197
Leidy, jr., J. W.
Reversed-phase high-performance liquid chromatographic purification of ^{125}I-labeled rat growth hormone-releasing hormone for radioimmunoassay 253
——, see Euerby, M. R. 239
Leyden, D. E., see Bogart, G. R. 209
Li, B., see Jin, Y. 394
Li, X.-N., see Kim, L.-S. 359
Lockley, W. J. S.
Regiochemical differences in the isotopic fractionation of deuterated benzoic acid isotopomers by reversed-phase high-performance liquid chromatography 413
Mangia, A., see Casoli, A. 443
Marsman, J. H.
——, Panneman, H. J. and Beenackers, A. A. C. M.
Automatic on-line pressurized liquid injection in gas chromatography 111
Mawhinney, T. P., see Landrum, D. C. 21
Meisch, H.-U., see Müller, G. 145
Miercke, L. J. W.
——, Stroud, R. M. and Dratz, E. A.
Preparative purification of functional bacteriorhodopsin by high-performance size-exclusion chromatography 331
Mucha, I.
—— and Tóth, G.
Separation of ^{125}I-labeled prostanoid derivatives by reversed-phase high-performance liquid chromatography 419
Müller, G.
—— and Meisch, H.-U.
Separation of inorganic anions by ion chromatography on dye-coated silica gel RP-18 145
Murray, S., see Williams, J. A. 454
Nakatani, S., see Kato, Y. 401
Newland, P.
——, Bingham, B., Tarelli, E. and Thomas, A. H.
High-performance gel permeation chromatography of meningococcal polysaccharides 406
Nielander, H. B., see Schotman, P. 301
Oi, T., see Kakihana, H. 179
Otani, M.
——, Umezawa, C. and Sano, K.
High-performance liquid chromatographic determination of pyrroloquinoline quinone with electrochemical detection and its application to bacterial samples 448
Panneman, H. J., see Marsman, J. H. 111
Partridge, L. Z., see Euerby, M. R. 239
Paterson, R. R. M.
—— and Kemmelmeier, C.
Gradient high-performance liquid chromatography using alkylphenone retention indices of insecticidal extracts of penicillium strains 153
Pennings, E. J. M.
——, Verpoorte, R., Goddijn, O. J. M. and Hoge, J. H. C.
Purification of tryptophan decarboxylase from a *Catharanthus roseus* cell suspension culture 311
Predieri, G., see Casoli, A. 443
Price, G. J.
—— and Dent, M. R.
Prediction of retention in gas–liquid chromatography using the UNIFAC group contribution method. I. Low-molecular-weight stationary phases 1

Reglero, G.
——, Herraiz, M., Herraiz, T. and Sanz, J.
Capillary gas chromatographic determination of volatiles in solid matrices by direct introduction using a programmable-temperature vaporizer 43
Rozen, A. J. van, see Schotman, P. 301
Salman, S. R.
——, Kamounah, F. S. and Tameesh, A. H. H.
Correlation between chromatography and ^1H NMR parameters in sterically substituted phenols 390
Sanagi, M. M., see Smith, R. M. 51
Sano, K., see Otani, M. 448
Sanz, J., see Reglero, G. 43
Sappa, E., see Casoli, A. 443
Schafer, W., see Bogart, G. R. 209
Schneider, G.
——, Kramell, R. and Brückner, C.
Separation of diastereomeric amino acid conjugates of jasmonic acid 459
Schotman, P.
——, Nielander, H. B., Van Rozen, A. J., Frankena, H., Schrama, L. H. and Gispen, W. H.
Microheterogeneity of the growth-associated neuronal protein B-50. Contribution of phosphorylation by protein kinase C 301
Schrama, L. H., see Schotman, P. 301
Secor, H. V., see Seeman, J. I. 169
Seeman, J. I.
——, Secor, H. V., Armstrong, D. W., Ward, K. D. and Ward, T. J.
Separation of homologous and isomeric alkaloids related to nicotine on a β-cyclodextrin-bonded phase 169
Smart, A. U., see Smith, J. R. L. 341
Smith, D. F.
——, Kleindienst, T. E. and Hudgens, E. E.
Improved high-performance liquid chromatographic method for artifact-free measurements of aldehydes in the presence of ozone using 2,4-dinitrophenylhydrazine 431
Smith, J. A., see Ford, J. C. 131
Smith, J. R. L.
——, Smart, A. U., Hancock, F. E. and Twigg, M. V.
High-performance liquid chromatographic determination of low levels of primary and secondary amines in aqueous media via derivatisation with 1,2-naphthoquinone-4-sulphonate 341
Smith, K. C., see White, R. L. 437
Smith, R. M.
—— and Sanagi, M. M.
Packed-column supercritical fluid chromatography of benzodiazepines 51
Snyder, L. R.
——, see Cox, G. B. 95
—— and Cox, G. B.
Preparative high-performance liquid chromatography under isocratic conditions. II. The role of column variables 85
——, Dolan, J. W. and Cox, G. B.
Preparative high-performance liquid chromatography under isocratic conditions. I. Craig simulations for heavily overloaded separations 63
Stroud, R. M., see Miercke, L. J. W. 331
Stouten, P., see Haasnoot, W. 319
Tambe, A. S., see Ayyangar, N. R. 33
Tameesh, A. H. H., see Salman, S. R. 390
Tarelli, E., see Newland, P. 406
Techel, D.
—— and Braumann, T.
Determination of cyclic nucleotide phosphodiesterase activity in cellular systems by ion-pair reversed-phase liquid chromatography 427
Thomas, A. H., see Newland, P. 406
Tian, Z.-Q., see Kim, L.-S. 359
Tóth, G., see Mucha, I. 419
Twigg, M. V., see Smith, J. R. L. 341
Umezawa, C., see Otani, M. 448
Van Rozen, A. J., see Schotman, P. 301
Venema, D. P., see Haasnoot, W. 319
Verpoorte, R., see Pennings, E. J. M. 311
Vijayalakshmi, M. A., see Kirchberger, J. 289
Vukmanic, D.
—— and Chiba, M.
Effect of organic solvents in sample solutions and injection volumes on chromatographic peak profiles of analytes in reversed-phase high-performance liquid chromatography 189
Wade, T. M., see Bogart, G. R. 209
Wang, J., see Kim, L.-S. 359
Wang, N.-H. L., see Whitley, R. D. 263
Wang, W.-K.
—— and Huang, S.-D.
Rapid determination of seven herbicides in water or isooctane using C_{18} and Florisil Sep-Pak cartridges and gas chromatography with electron-capture detection 121
Ward, K. D., see Seeman, J. I. 169
Ward, T. J., see Seeman, J. I. 169
White, R. L.
——, Demarco, A. C. and Smith, K. C.
Analysis of o-phthalaldehyde derivatives in fermentation broths by high-performance liquid chromatography 437

Whitley, R. D.
——, Brown, J. M., Karajgikar, N. P. and Wang, N.-H. L.
Determination of ion-exchange equilibrium parameters of amino acid and protein systems by an impulse response technique 263

Williams, J. A.
——, Davies, D. S. and Murray, S.
Synthesis of 7α-thiohydroxycholesterol involving the resolution of epimeric thiocholesterols by high-performance liquid chromatography 454
Zhang, Y., see Kim, L.-S. 359